



National Diet and Nutrition Survey Years 1-4 2008/09-2011/12

User Guide for UK Data (*core & country boost data*)



At **NatCen Social Research** we believe that social research has the power to make life better. By really understanding the complexity of people's lives and what they think about the issues that affect them, we give the public a powerful and influential role in shaping decisions and services that can make a difference to everyone. And as an independent, not for profit organisation we're able to put all our time and energy into delivering social research that works for society.

The Medical Research Council Elsie Widdowson Laboratory (MRC EWL), formerly called MRC Human Nutrition Research (HNR), conducts internationally-renowned research aimed at improving health through nutrition. Our mission is to conduct nutrition research and surveillance to improve the health of the population and to inform nutrition policy and practice.

University College London (UCL) is an outstanding institution, recognised as one of the world's most advanced universities and valued highly by its community of staff, students, alumni, donors and partners and by the wider community. It provides an outstanding education to students from across the globe that imparts the knowledge, wisdom and skills needed by them to thrive as global citizens and is committed to achieving maximum positive social, environmental and economic benefit through its achievements in education, scholarship, research, discovery and collaboration.

NatCen Social Research
35 Northampton Square
London EC1V 0AX
www.natcen.ac.uk

A Company Limited by Guarantee
Registered in England No.4392418.
A Charity registered in England and Wales (1091768) and Scotland (SC038454)

Contents

Glossary 5

1	Introduction	8
1.1	The National Diet and Nutrition Survey Rolling Programme (NDNS RP) Years 1-4 User Guide	8
1.2	The NDNS RP	8
1.3	NDNS RP and sodium study reports	9
2	Survey design	11
2.1	Comparison study	12
2.2	Fieldwork	12
2.3	Survey structure	12
3	Archive documentation	14
4	Using the data	15
4.1	Differences from previous versions of the NDNS RP Y1-4 datasets	15
4.2	Years 1-4 datasets	16
4.3	UK Nutrient Databank	17
4.4	Variables in the datasets	18
4.5	Missing values conventions	18
4.6	Merging datasets	19
4.7	Serial number composition	19
5	Using the dietary data files	21
6	Using doubly labelled water (DLW) data.....	24
7	Using 24-hour urine data.....	25
8	Using blood data.....	27
9	Weighting variables	29
9.1	Description of weights	29
9.2	Single weights for all individuals	30
9.3	The Year 1 Run-in sample.....	30
9.4	Individual non-response weight.....	30
9.5	Nurse interview non-response weight.....	31
9.6	RPAQ non-response weight	32

9.7	ActiGraph non-response weight	32
9.8	Blood sample non-response weights	33
9.9	24-hour urine sample non-response weights	33
9.10	Selecting the appropriate weight variable.....	35

Appendix A UK Nutrient Databank..... 37

A.1	Introduction to the UK Nutrient Databank.....	37
A.2	Description of the Nutrient Databank fields and variables	38
A.2.1	Food name and food code	38
A.2.2	Food code markers	38
A.2.3	Sub food group name and code	38
A.2.4	Water and vitamin loss	38
A.2.5	Maximum weight	39
A.2.6	Base and unit (F codes only)	39
A.2.7	Dilution	39
A.2.8	Edible portion	39
A.2.9	Comments and descriptions	39
A.2.10	Nutrition data.....	40
A.3	Quality assurance	40
A.4	Changes to the databank between Years 1 to 4	40

Appendix B Doubly labelled water (DLW) 43

B.1	Total Body Water (L)	43
B.2	Lean Body Mass (kg) and Fat Mass (kg)	43
B.3	Basal Metabolic Rate (according to Schofield's equations, using age, sex, height and weight).....	44
B.4	Total Energy Expenditure (kJ) (calculated according to Schoeller's methods) 44	
B.5	Activity energy expenditure (AEE) and physical activity energy expenditure (PAEE)	45
B.6	Physical activity level (PAL)	45

Glossary

25-OHD	25-hydroxyvitamin D
5-A-Day	5 portions of fruit and vegetables per day (these are 5 x 80g portions)
AEE	Activity Energy Expenditure
BMR	Basal Metabolic Rate
Ca	Calcium
CAPI	Computer Assisted Personal Interview
CO ₂	Carbon Dioxide
Databank	UK Nutrient Databank
DH	Department of Health
DLW	Doubly Labelled Water
Fe	Iron
FM	Fat Mass
FQ	food quotient
FSA	Food Standards Agency
G	Gram
GOR	Government Office Region
H	Hydrogen
HbA1c	Glycosylated Haemoglobin or Haemoglobin A1c
HRP	Household Reference Person
IMS	Intrinsic Milk Sugars
Kcal	kilocalorie
Kg	Kilogram
KJ	Kilojoule
L	Litre

LBM	Lean Body Mass
LIDNS	Low Income Diet and Nutrition Survey
LDL	Low Density Lipoprotein Cholesterol
LRNI	Lower Recommended Nutrient Intake
µg	Microgram
MAFF	Ministry of Agriculture, Fisheries and Farming
MG	Milligram
Mins	Minerals
MFP	Main Food Provider
MRC	Medical Research Council
MRC EWL	Medical Research Council Elsie Widdowson Laboratory
MRC HNR	Medical Research Council Human Nutrition Research
NatCen	NatCen Social Research
NDNS	National Diet and Nutrition Survey
NISRA	Northern Ireland Statistics and Research Agency
NMES	Non-Milk Extrinsic Sugars
NSP	Non-Starch Polysaccharide
O	Oxygen
PABA	Para-aminobenzoic acid
PAEE	Physical Activity Energy Expenditure
PAF	Postcode Address File
PAL	Physical Activity Level
PHE	Public Health England
PSU	Primary Sampling Unit
RNI	Reference Nutrient Intake

RP	Rolling Programme
RPAQ	Recent Physical Activity Questionnaire
Se	Selenium
sTfR	Soluble Transferrin Receptors
Sub	Subsidiary
TBW	Total Body Water
TEE	Total Energy Expenditure
Trig	Triglyceride or Triacylglycerol
UCL	University College London
UK	United Kingdom
Vit C	Vitamin C
Vits	Vitamins
Yr1-4	Years 1 to 4
Zn	Zinc

1 Introduction

1.1 The National Diet and Nutrition Survey Rolling Programme (NDNS RP) Years 1-4 User Guide

This is the User Guide for the NDNS RP Years 1-4 dataset; the document supersedes the previous version (available on the UK Data Archive until **October 2016**). Section 4.1 of this Guide provides detail about the updates that have been made to the NDNS RP Years 1-4 datasets to correct discrepancies and/or to provide additional and updated data.

Separate User Guides will be produced for the Years 5-6 and subsequent datasets.

1.2 The NDNS RP

The NDNS RP is designed to assess the diet, nutrient intake and nutritional status of the general population aged 1.5 years and over living in private households in the UK. The NDNS RP is jointly funded by Public Health England (PHE),¹ an executive agency of the Department of Health, and the UK Food Standards Agency (FSA)^{2,3,4} and for this survey period was carried out by a consortium of three organisations: NatCen Social Research (NatCen),⁵ MRC Elsie Widdowson laboratory (MRC EWL), formerly known as MRC Human Nutrition Research (HNR),⁶ and the University College London Medical School (UCL).⁷ Fieldwork in Northern Ireland is carried out by the Northern Ireland Statistics and Research Agency (NISRA).⁸

The NDNS RP provides the only source of high quality nationally representative data on the types and quantities of foods consumed by individuals, from which estimates of nutrient intake for the population are derived.⁹ Results are used by Government to develop policy and monitor progress on diet and nutrition objectives of UK Health Departments, for example those set out in the Healthy Lives, Healthy People White Paper in England.¹⁰ The food consumption data are also used by FSA to assess exposure to chemicals in food, as part of the risk assessment and communication process in response to a food emergency or to inform negotiations on setting regulatory limits for contaminants.

The NDNS programme began in 1992 as a series of cross-sectional surveys, each covering a different age group. Since 2008, the NDNS has been a RP covering adults and children aged 1.5 years and over. The NDNS collection also includes separate survey assessments of dietary sodium in adults.

The specific aims of the NDNS RP are to:

- provide quantitative data on the food and nutrient intakes, sources of nutrients and nutritional status of the UK population aged 1.5 years and above;
- provide information on trends in food consumption, nutrient intake and nutritional status in different age groups;
- describe the characteristics of individuals with intakes of specific nutrients above or below the national average;
- produce a database of food consumption which will be used to calculate intakes of natural toxicants, contaminants, additives and other food chemicals;
- measure blood and urine indices that provide evidence of nutritional status or dietary biomarkers, and to relate these to dietary, physiological and socio-demographic data;
- provide height, weight and other anthropometric measurements and examine their relationship to socio-demographic, dietary, biochemical and health data;
- monitor the diet of the population to establish the extent to which it is adequately nutritious and varied;
- monitor the extent to which the diets of population sub-groups vary from expert recommendations;
- assess total energy expenditure and physical activity levels and patterns in the study population.

1.3 NDNS RP and sodium study reports

Further information about the NDNS collection can be found on the gov.uk site: <https://www.gov.uk/government/collections/national-diet-and-nutrition-survey>

Detail about Years 1 to 4 of the NDNS RP are provided in the published NDNS RP Years 1 to 4 UK and devolved country reports:

UK report: <https://www.gov.uk/government/publications/national-diet-and-nutrition-survey-results-from-years-1-to-4-combined-of-the-rolling-programme-for-2008-and-2009-to-2011-and-2012>

National Diet and Nutrition Survey Rolling Programme (NDNS RP) Years 1-4 User Guide

Scotland report: <http://www.foodstandards.gov.scot/national-diet-and-nutrition-survey-rolling-programme-results-years-1-4-combined-scotland-200809>

Northern Ireland report: <https://www.food.gov.uk/northern-ireland/researchni/ndns-ni>

Wales report: <http://gov.wales/statistics-and-research/national-diet-nutrition-survey-rolling-programme/?lang=en>¹

Folate report: <https://www.gov.uk/government/statistics/national-diet-and-nutrition-survey-supplementary-report-blood-folate>

Further information about the England 2011 sodium study is provided in the published report:

<https://www.gov.uk/government/publications/assessment-of-dietary-sodium-levels-among-adults-aged-19-64-in-england-2011>

¹ Note that the Wales report covers Years 2 to 5. Data for Year 5 will be included in the Years 5/6 dataset.

2 Survey design

The NDNS RP is a survey of the food consumption, nutrient intakes and nutritional status of people aged 1.5 years and older living in private households. The survey is carried out in all four countries of the UK and is designed to be representative of the UK population.

The sample was drawn from the Postcode Address File (PAF), a list of all the addresses in the UK. In order to improve cost effectiveness the addresses were clustered into Primary Sampling Units (PSUs), small geographical areas based on postcode sectors, randomly selected from across the UK. A list of addresses was randomly selected from each PSU.

The survey aimed to collect data from a UK representative sample of 1000 people per year, 500 adults (aged 19 years and older) and 500 children (aged 1.5 to 18 years). In addition, additional recruitment was carried out in the devolved countries to increase sample sizes (the aim was to increase the sample size to 400 individuals (200 adults, 200 children) per year in Scotland and 200 per year (100 adults, 100 children) in Wales and in Northern Ireland.

Overall for Years 1 to 4 combined a UK sample of 21,573 addresses was selected from 799 PSUs. A total of 27 addresses were randomly selected in each selected PSU. At each address, the interviewer established the number of households and, in cases where there were two or more, selected one household at random. In order to achieve (as far as possible) equal numbers of adults and children in the sample, at some addresses only children were selected to take part. The 27 addresses were randomly allocated to one of two groups to determine whether an adult and a child, or a child only, was selected for interview.

For Years 1 to 3, at nine of the selected addresses the interviewer selected one adult (aged 19 years and over) and, where present, one child (aged 1.5 to 18 years) for inclusion in the survey (basic addresses). The remaining 18 addresses were for a “child boost” where the interviewer only carried out interviews in households with children (aged 1.5 to 18 years). In Year 4, quarters 1 and 2, this split of addresses changed to 11 basic addresses and 16 child boost. In quarter 3 and 4 this split was 10 basic addresses and 17 child boost addresses.

For households containing more than one eligible person (adult and/or child), interviewers selected the participant(s) using a random selection procedure.

In total 3,450 adults and 3,378 children gave fully productive interviews (consisting of three or four diary days). Overall in Years 1 to 4 combined, 51% of adults (1,769) and 27% of children (902) who had completed a diary went on to give a blood sample. Sixty per cent of adults (2,074) and

58% of children aged 4 to 18 years (1,602) who completed a diary agreed to provide a 24-hour urine sample.

2.1 Comparison study

Prior to the launch of the first year of main stage fieldwork, a comparison study of two alternative dietary assessment methods (randomly allocated to sampled addresses) was carried out in 2007. Over 1,100 adults and children took part with around half participating in interviewer-administered 24-hour dietary recalls (repeated on four non-consecutive days) and the others keeping a four-day estimated (unweighed) food diary on consecutive days. The NDNS Project Board considered the findings and decided that the four-day estimated diary (hereafter referred to as the “four-day food diary”) should be used for the RP.

2.2 Fieldwork

Field work was conducted throughout the year (from February 2008 through August 2012) in order to take into account potential seasonal variations in food consumption. February and March 2008 points were part of the ‘Run-In’ to Year 1 where protocols and procedures were tested before the main stage fieldwork period launched in April 2008. It consisted of ten PSUs issued over two months and was carried out in all four UK countries. The Run-in sample was drawn in the same way as the NDNS RP main stage core sample and fieldworkers followed the same protocols and procedures as in the main stage. The Run-in data have therefore been combined with Year 1 of the main stage data and participants from the Run-in sample had a serial ID starting with ‘9’, see chapter 9, section 9.3 of this user guide regarding Run In weightings information.

2.3 Survey structure

There are two main parts to the survey: an interviewer stage and a nurse visit.

Stage 1: Interviewer visit:

- Four-day food diary²
- Face-to-face Computer Assisted Personal interview (CAPI)³

² See Chapter 5 of the UK Years 1 to 4 report and Chapter 6 of this User Guide for more details.

-
- Height and weight measurements
 - Smoking and drinking self-completion questionnaires
 - Physical activity self-completion questionnaire or ActiGraph

On successful completion of the interviewer stage (including three or four completed days of the food diary), each participant was invited to take part in the next stage, a visit from a nurse.

Stage 2: Nurse visit:

- Fasting blood sample (aged four years and over)
- Non-fasting blood sample (aged 1.5 to three years)
- 24-hour urine collection (aged four years and over)
- Physical measurements (e.g. waist and hip circumference)
- Blood pressure
- Collection of information about prescribed medicines & dietary supplements

In addition, a subsample of participants in Years 1 and 3 were recruited for a Doubly Labelled Water (DLW) sub-study to measure energy expenditure (see Chapter 7 of this User Guide).

³ Collecting information on shopping and food preparation practices, cooking facilities in the household, eating habits and food avoidance. Also included a section for the Main Food Provider (MFP).

3 Archive documentation

The NDNS RP documentation on the UK Data Archive has been organised into the following sections:

- **Survey documents**

This contains the CAPI documentation for the interviewer visit and nurse schedules, self-completion questionnaires, showcards and consent forms.

- **Data related documents**

NDNS RP Yr1-4 Variable List - this contains a list of the variables on each dataset and the survey year to which it applies

NDNS RP Yr1-4 Derived Variables – this contains the SPSS syntax specification for each of the derived variables included in the data.

- **Supporting documents**

This contains details of food coding and instructions for office editing of the diaries and coding of the CAPI data and self-completions. Also, documents related to four-day food diary, interviewer and nurse project instructions and protocols.

Note that the questionnaires show the variable names used in the CAPI programme for Year 4. In some cases the variables in the data set have a different name where changes have been made to the question or routing between years. In these cases consult the questionnaire for each individual survey year which can be accessed via the UK Data Archive.

4 Using the data

4.1 Differences from previous versions of the NDNS RP Y1-4 datasets

A number of updates, described below, have been made to the NDNS RP Y1-4 datasets to correct discrepancies and/or to provide additional and updated data. The individual datasets are listed at 4.2 below. Descriptive statistics calculated using variables in the revised datasets may therefore differ from descriptive statistics performed on data obtained from previous versions of these datasets due to amendments made to variables where applicable. A summary of the variables that have been updated is provided below and further details can be found in the relevant chapters of this User Guide:

5-A-Day variables have been amended to exclude the food groups listed in Appendix A (Chapter 5) of the NDNS RP Years 1-4 published reports).

Data has been amended for the DLW sub-study which forms part of the NDNS RP Y1-4, to correct a number of minor discrepancies in height, weight and dose weight, and one discrepancy in TEE calculation. All discrepancies had very small impacts on: FM_KG; BMR_KJ_DAY_1; TEE; TEEREP_KJ_DAY_1; CV2 and PAL (Chapter 6). Food quotient (FQ) values have been updated to take account of small changes in dietary intake assessment methods over time. This has an impact on the variables: RQ; TEEREP_KJ_DAY_1 and PAL (chapter 6 of this User Guide).

Additional urinary sodium variables have been added to the NDNS RP Years 1-4 dataset to provide adjusted 24-hour sodium excretion data, following multiplication by a method-specific factor, for results previously reported as part of the NDNS RP reports (see Chapter 7 of respective published reports for more information). The adjustment factor was introduced at the time of reporting results from the 2014 England urinary sodium survey in order to improve accuracy and to enable comparability across different survey assessments.

Years 1-4 data for Fat Soluble Vitamins have been amended by applying an analyte specific conversion factor to enable comparability with the method used from Year 5 onwards (see Chapter 8 of this User Guide).

High 25-OHD concentrations compromised by sample-specific interference in the old-formulation of Liaison kits have been replaced by 25-OHD concentrations obtained using the new-formulation Liaison kits (Chapter 8 of this User Guide and Chapter 6 of the NDNS RP published reports).

25-OHD concentrations have been retrospectively standardised against an in-house MRC EWL (then called MRC HNR) LCMS/MS method; the standardised concentrations are archived in addition to the unadjusted data (Chapter 8 of this User Guide).

Data have been added for folate variables which have been presented in the NDNS RP supplementary folate report. The variable listing has been updated to reflect all changes and provides a complete list of all of the variables included in the NDNS RP UK Years 1 to 4 datasets (Chapter 8 of this User Guide).

Data have been added for the weights used in the country-specific reports. The variable listing has been updated to reflect all changes and provides a complete list of all of the variables included in the NDNS RP UK Years 1 to 4 datasets (Chapter 9 of this User Guide).

4.2 Years 1-4 datasets

Data collected during the survey are contained in different data files described below.

Name of Dataset	No. of records	Description of Dataset
NDNS_Yr1-4a_indiv_cere	6,828	Contains data for all fully productive individuals i.e. completed three/four food diary days. It contains information from the household questionnaire, main individual schedule, self-completions, physical measurements and nurse visit (where one occurred). It also includes blood and 24-hour urine sample results and data from the DLW sub-study including energy expenditure data for relevant participants.
NDNS_Yr1-4a_hhold_cere	17,016	Contains data on household composition, sex, age and marital status for all individuals in co-operating households.

NDNS_Yr1-4a_FoodLevelDietaryData__core	619,173	Diary data. Includes nutrient data and disaggregation at food level. Also, shows who else was present at the eating occasion, where the participant was located, whether the television was on and whether or not the participant was sitting at a table.
NDNS_Yr1-4a_DayLevelDietaryData_Foods__core	27,183	Daily food consumption data calculated using recipe main food groups and recipe sub food groups data.
NDNS_Yr1-4a_DayLevelDietaryData_Nutrients__core	27,183	Daily intakes of macronutrients, micronutrients and disaggregated foods.
NDNS_Yr1-4a_PersonLevelDietaryData__core	6,828	Mean intakes of nutrients, food consumption data calculated using recipe main food groups and recipe sub food groups data plus disaggregated food at the participant level. Also includes derived variables such as LRNI and RNI indicators and percentages.

4.3 UK Nutrient Databank

The UK nutrient databank used in the NDNS RP is saved as four different data files, one for each survey year. Details about the nutrient databank and its use can be found in Appendix A of this User Guide.

Name of Dataset	Description of Dataset
Year1DataBank_2014-06-13	Contains nutrient data assigned to foods and supplements for Year 1
Year2DataBank_2014-06-13	Contains nutrient data assigned to foods and supplements for Year 2
Year3DataBank_2014-06-13	Contains nutrient data assigned to foods and supplements for Year 3
Year4DataBank_2014-06-13	Contains nutrient data assigned to foods and supplements for Year 4

4.4 Variables in the datasets

The individual and household datasets contain questionnaire variables (excluding variables used for administrative purposes), demographic information including household composition, laboratory results and derived variables.

The dietary datasets contain variables coded from the diaries at food, day and person levels, plus dietary reference values and derived variables.

The variables included in all the datasets are detailed in the “**NDNS RP Yr1-4 List of Variables**” document in the data section of the documentation. This document is the best place to look in order to plan your analysis. It includes:

- Major categories of variables (e.g. General Health, Blood Sample, Day Level Dietary Nutrients)
- Sub categories of variables (e.g. Longstanding illness (within General Health), Measurements from laboratory analysis (within the Blood Sample, Urine Collection and DLW sample sections), Nutrients including supplements (within Day Level Dietary Nutrients)
- Source of each variable (e.g. Individual questionnaire, Diary, Nurse visit, Self-completion booklet, Laboratory, Derived variable etc.)

Details of the question wording relating to a variable in the household and individual datasets is provided in the interview section documentation for the appropriate survey year (all variables in the dataset are given by name in the copy of the interview schedules provided on the UK Data Archive). The “**NDNS RP Yr1-4 Derived Variables**” document on the UK Data Archive provides information on how the variables were derived.

4.5 Missing values conventions

Missing value conventions are applied to most of the derived variables as well as the original questionnaire variables. The “**NDNS RP Yr1-4 Derived Variables**” specification should be consulted for details.

- 1 Not applicable. This code is used to signify that a particular variable did not apply to a given participant because of internal routing (e.g. questions for children only) or because the participant did not participate in a particular element of the survey (e.g. refused a nurse visit).

-
- 4 Question not applicable to survey year. This code indicates that that particular question was not asked in a survey year. For example the variable *Operat* – whether the respondent has any surgical procedures in the last 12 months, was only asked of Year 1 respondents and therefore all Year 2, Year 3 and Year 4 respondents will be coded as a -4.
 - 8 Don't know/Can't say
 - 9 No answer/Refusal

The above conventions apply to the majority of the variables on the data, however some variables have been attributed specific missing value codes, for example blood, urine and DLW results on the individual dataset, and “5-A-Day” variables on the dietary data files. The description for each of the missing value codes are specified in the each variable value label.

For a full list of variables on the dataset and to which survey they apply see the “**NDNS RP Yr1-4 List of Variables**”.

4.6 Merging datasets

As various data are contained in different datasets, users may need to merge several datasets together for the purposes of their analysis. Individual serial number, survey year, age, sex and country variables are included in all the datasets for consistency.

4.7 Serial number composition

Serial numbers on the data consist of the following:

Variable name	Definition	Composition	File/files included
SERIALH	Household serial number	7 digits. The same number is allocated to each member of the same household. The first number corresponds to the survey year ¹¹	Household and individual files
SERIALP	Individual identifier for each household member in a productive household	9 digits. SERIALH + PGRID	Household file only
SERIALI	Individual serial number for each productive individual (i.e. completed 3 or more diary days	8 digits SERIALH + ADCHILD ADCHILD is coded 1 for adult and 2 for child	Household, Individual and all Dietary files

The individual file also contains the person number of the Household Reference Person (HRP) and the Main Food Provider (MFP) (variables HRPNO and MFPNUM respectively). To create individual serial numbers for either the HRP or MFP, add HRPNO or MFPNUM to SERIALH.

Note that the HRP or MFP numbers correspond to the person number within each household. Therefore, due to the recoding of each productive individual to a 1 for the adult and 2 for the child, the HRP or MFP may not be the same individual although they may have the same serial number and vice versa. An example is show below for a household consisting of 2 adults and 2 children.

	SERIALH	PGRID	SERIALP	Productive Y/N	SERIALI	Is HRP? Y/N	Is MFP? Y/N	HRP serial	MFP serial
Adult 1	1010101	1	101010101	No	N/A	No	Yes	N/A	10101011
Adult 2	1010101	2	101010102	Yes	10101011	Yes	No	10101012	N/A
Child 1	1010101	3	101010103	No	N/A	N/A	N/A	N/A	N/A
Child 2	1010101	4	101010104	Yes	10101012	N/A	N/A	N/A	N/A

In this example, the MFP and the individual are not the same person even though they have the same MFP and individual serial numbers. For clarification and cross checking, the age and sex of the HRP and MFP have been provided.

5 Using the dietary data files

It is important to note the following when using the dietary data from Year 1 to 4 of the NDNS RP:

Days of the week

- Year 1 of the RP was designed to include two week days and two weekend days for each person. As a result some bias may have been introduced into the results for foods and nutrients for which intakes differ between week days and weekend days, for example alcohol.
- To redress the over-representation of weekend days and non-representation of Wednesdays in Year 1, the food diary recording period was changed from Year 2 onwards so that all days of the week would (as far as possible) be equally represented.
- Years 3 and 4 were designed so that all days of the week were evenly represented. However, in the Years 1 to 4 combined data, there remains a slightly higher proportion of weekend days than weekdays.

Dietary coding

- Changes have been made to the dietary coding of homemade recipes and some purchased convenience foods in the NDNS RP, compared with previous standalone NDNS, which should be considered by those wishing to calculate food consumption data. In the RP all individual ingredients of a homemade recipe as reported in the food diary, or components of the purchased product as described on the food packaging, have been coded as their separate food codes and linked together under the appropriate Recipe Food Group, which highlights that those food codes were consumed together in one composite dish. The following variables should be used when calculating food consumption data;
 - RECIPEMAINFOODGROUPCODE
 - RECIPEMAINFOODGROUPDESC
 - RECIPESUBFOODGROUPCODE
 - RECIPESUBFOODGROUPDESC
- An example is provided here: A homemade dish of Thai chicken curry containing chicken, Thai curry sauce, and onion would appear in the RP Food Level dietary dataset as three entries with the food names; CHICKEN BOILED LIGHT MEAT ONLY, THAI CURRY SAUCE PURCHASED, and ONIONS BOILED, linked to the MAINFOODGROUPDESC of

“chicken and turkey dishes”, “miscellaneous” and “vegetables not raw”, respectively. As these three foods were consumed together in one composite dish they are assigned to the RECIPESUBFOODGROUPDESC of Other chicken/turkey including homemade recipe dishes. In previous NDNS datasets this homemade dish would have shown as one entry, assigned as one food name, and the proportions of the composite dish consumed would not have been known.

- To estimate absolute food consumption of one specific food type examine the FOODNAME and MAINFOODGROUPDESC variables, whilst examining disaggregation variables of any foods that are composites (NB disaggregation data is only provided for certain categories of meat, fish, fruit and vegetables). For example, to estimate absolute intakes of sausages from all sources you would need to include all the specific discrete portions of sausages, as well as calculate the percentage of sausages within all composite foods such as meat pies.
- All foods consumed have a base unit of grams that is, the amount consumed is described in grams. The exceptions are dietary supplements and artificial sweeteners. These have a base unit based on their form i.e. tablet, teaspoon. To avoid errors when calculating consumption, these have only been included in the food level dietary data file. When using this file, it should be noted that, for dietary supplements and artificial sweeteners, the value in the Total_Grams column is not a value in grams but a value in terms of the base unit, i.e. 0.5 for a granulated artificial sweetener would refer to 0.5 of a teaspoon not 0.5 grams.

5-A-Day calculations

- Full details of the methods used for the disaggregation of meat, fish, fruit and vegetables are provided in Appendix A of the UK and devolved country NDNS RP Years 1 to 4 reports.
- Variables used for 5-A-Day are calculated according to methodology described in Appendix A (section A.4.3) of the NDNS RP published reports. Data were incorrect in the previous version of the dataset (available on the UKDA until **October** 2016) and in the 2014 published reports due to certain food groups not being appropriately excluded. Data generated using variables in the current dataset may therefore not match the variables in original versions of the NDNS RP Years 1 to 4 reports or previous downloads of these datasets as these food groups had not been excluded previously.

-
- For further details regarding all other dietary data methodologies used in the RP refer to Appendix A in the UK and devolved country NDNS RP Years 1 to 4 reports.

Comparing NDNS RP data to previous NDNS data

In comparing the NDNS RP data to past published reports of the NDNS, the differences in duration of assessment must be taken into account. Dietary assessment over a four day period will provide similar mean intakes from assessment over a seven day period, but the variation in intakes will be different as will the percentage of participants consuming any given food over the recording period (percentage consumers), especially for foods that are consumed infrequently. Moreover, estimates of proportions of individuals above or below certain cut-off values, such as Lower Reference Nutrient Intakes (LRNIs) will be affected by assessments of different duration. For this reason, for the previous NDNS of young people aged 4 to 18 years and NDNS of adults aged 19 to 64 years reports, the seven day information from previous NDNS surveys was converted to four days using bootstrapping techniques. Details of the bootstrapping methodology used and descriptive statistics of bootstrapped data for NDNS foodgroups are provided in Appendix K of the NDNS RP Years 1 to 4 report (see section 1.3 of this User Guide for links to published reports).

6 Using doubly labelled water (DLW) data

It is important to note the following when using the DLW data. The variable “BMR (kJ)” is calculated according to Schofield’s equations and the variable “TEE (kJ)” is calculated from Schoeller’s equation. For further details and the equations used refer to Appendix B of this document.

The archived DLW data for the NDNS RP Years 1-4 has been updated, please see section 4.1 of this User Guide for further details.

7 Using 24-hour urine data

Urinary analyte data (including urinary sodium data used to estimate salt intake) have been supplied for fully continent participants over the age of four years who were deemed to have provided a complete sample. For those who provided a full 24-hour urine collection, but urinary analyte results could not be obtained, the interview data have been included in the dataset but these respondents have been attributed a specific missing value code for each result variable and given a urine weighting of '0'.

Urine collections for children aged 4 to 6 years and 7 to 10 years were classified as 'complete' using two different sets of criteria to determine which samples are complete; 'standard criteria' ('complete by PABA' or 'complete by claim')¹² or 'claim only' where collections were regarded as 'complete' when they were claimed to include all urine passed for 23-25 hours from the start time *irrespective* of PABA excretion.¹³ This approach reflects the uncertainty in ascertaining completeness of collections in this age group (more details are provided in Chapter 7 and Appendix T of the NDNS RP UK and devolved country Years 1 to 4 reports). Urine collections for children aged 11 to 18 years and adults aged 19 years and over were classified as 'complete' by 'standard criteria' ('complete by PABA' or 'complete by claim').¹² The 24-hour urinary analyte excretion results and the weight of the 24-hour urine collection (in kg) have only been provided for complete collections, but concentrations of all urinary analytes are provided for all urine collections irrespective of completeness.

Measurement of urinary sodium was carried out using an ion selective electrode (ISE) on the Siemens Dimension® Xpand clinical chemistry system with the QuikLYTE® module. During the time that these urine samples were assayed this analytical method showed slight negative bias for urinary sodium measurements relative to inter-laboratory consensus measurements. A method-specific correction factor (1.052) was derived from comparison studies against an instrument which provided accurate results. Corrected results for urinary sodium excretion have now been provided in addition to the original results in the current NDNS dataset.

Data for the following urinary variables (multiplied by a method-specific factor) have been added to the datasets to enhance accuracy and enable comparison with previous urinary sodium survey data obtained with different methods:

Na_mmol_Corrected	Urinary sodium concentration (mmol/L) – Adjusted by method-specific correction factor (1.052).
-------------------	--

Na_mmol_24h_ALL_Corrected	Urinary sodium excretion (mmol/24-hours) - Results for all urine collections deemed complete. Calculated using corrected sodium concentration.
Na_mmol_24h_STANDARD_Corrected	Urinary sodium excretion (mmol/24-hours) - Results for urine collections deemed complete EXCEPT where ONLY completeness criteria is aged 4-10, 0 missed samples, collection for between 23 to 25 hours. Calculated using corrected sodium concentration.
Na_mmol_24h_4_10CLAIM_Corrected	Urinary sodium excretion (mmol/24-hours) - Results for urine collections deemed complete for participants aged 4-10 by claim only (not complete by PABA). Calculated using corrected sodium concentration.

8 Using blood data

The complete blood results dataset have been supplied for those aged 1.5 years and over including both data for blood analytes published in the NDNS RP Years 1 to 4 report as well as data for additional blood analytes measured but not reported. For those who provided a blood sample but results could not be obtained, the interview data has been included in the dataset but these respondents have been attributed a specific missing value code for each blood result variable and given a blood weight of 0.

Low Density Lipoproteins (LDL), Triglycerides (Trig) and Glucose (Glucose) result data are only provided for fasted blood samples. Glucose (Glucose), Homocysteine (Homocysteine), Soluble Transferrin Receptor (sTfR), Selenium (Se) and Zinc (Zn) result data has been supplied for those aged 7 years and over.

Glucose and Haemoglobin A1c (HbA1c) were funded separately by Diabetes UK and were measured in blood samples collected for Years 1 to 4 of the NDNS RP. Thyroid function hormones were funded separately by Addenbrooke's hospital, Cambridge (UK) and were measured in blood samples collected for Years 2 to 3 of the NDNS RP. These data are included in the archive dataset.

Fat soluble vitamins

From Year 5 onwards there was a change in the method used for the analysis of fat soluble vitamins. The data in the archived datasets have had the following conversion factors applied so that the data now match data generated by the new method presented in the Wales Years 2 to 5 report and subsequent NDNS RP reports. The original values presented in the NDNS RP Years 1 to 4 reports can be obtained by dividing the data by the conversion factor (provided in the table below) for the analyte in question:

Analyte	Conversion factor
Retinol	0.84
α -tocopherol	0.82
γ -tocopherol	0.64
Lutein and zeaxanthin	0.70
α -cryptoxanthin	0.69

β -cryptoxanthin	0.83
Lycopene	1.25
α -carotene	0.83
β -carotene	0.88

Vitamin D status (25-hydroxyvitamin D)

25-hydroxyvitamin D (25-OHD) was measured using the Diasorin Liaison method. These kits were later withdrawn and reformulated by the manufacturers because some samples gave inaccurately high results, greater than 100nmol/L, as a result of sample-specific interference. Fifty-five potentially-affected samples (results >100nmol/L) from Years 1-4 were reanalysed using the reformulated kits after the publication of the Years 1 to 4 report and the current dataset includes the reissued results. There are therefore fewer 25-OHD results >100nmol/L in the current dataset.

Both Diasorin Liaison 25-OHD methods were standardised against an in-house MRC EWL (then called MRC HNR) LCMS/MS method which gave results equivalent to the international gold-standard isotope-dilution-liquid chromatography-tandem mass spectrometry methods, as validated under the Vitamin D Standardization Program (VDSP). Concentrations after standardisation are approximately 3% higher than the results as assayed. Standardised results (v25OHD_Std) are archived alongside the unadjusted data (v25OHD).

Folate

Year 1 to 4 core and boost data for red blood cell folate, serum total folate and folic acid have been included in the dataset (please refer to the variable listing).

9 Weighting variables

9.1 Description of weights

The NDNS RP requires weights to correct for differences in sample selection and response. The weights adjust for differential selection probabilities of households and individuals, non-response to the individual and RPAQ questionnaires, non-response to the nurse visit and non-response to the blood sample. Non-response weights were generated using a mixture of non-response modelling and calibration weighting methods. The sample has been drawn, and the weights designed, so that all four survey years can be used together. Seven weights were generated for the NDNS RP Years 1-4 dataset- these are described in Table 1.

Table 1 Description of the NDNS RP Years 1-4 weights

Weight name	Description of weight	Use for
wti_Y1234	Weight for non-response by adults and children to the individual questionnaire and diary	Any analysis of individuals using data from the individual questionnaire or diary. Including analysis of Smoking & Drinking data (collected in self-completions & CAPI)
wtn_Y1234	Weight for non-response by individuals to the nurse visit	Any analysis of individuals using data collected at the nurse visit
wtb_Y1234	Weight for non-response by individuals to the blood sample	Any analysis of individuals using blood sample data
wtr_Y1234	Weight for analysis of RPAQ (all individuals aged 16+)	Any analysis of RPAQ info for individuals aged 16+
wtu_Y1234v1	Weight for analysis of urine data	Any analysis of individuals using urine sample data (with completeness based on the 'standard adult criteria' for all individuals aged 4 years and over)
wtu_Y1234v2	Weight for analysis of urine data	Any analysis of individuals using urine sample data (with sample completeness based on complete by claim for individuals aged 4-10 years and the 'standard adult criteria' for individuals aged 11-15 years.)
wta_Y1234	Weight for analysis of ActiGraph data	Any analysis of individuals using ActiGraph data

9.2 Single weights for all individuals

Unlike previous datasets there is a single weight for all individuals, rather than separate weights for adults and children. This means the sample needs to be filtered by age to ensure the correct ages are included. However, this means different age breaks to those presented in the NDNS RP Years 1 to 4 reports can be used, i.e. 16 to 18 year olds can be combined with adults (19 years and over), which allows more flexibility in reporting.

9.3 The Year 1 Run-in sample

Prior to the launch of Year 1 fieldwork, the NDNS RP used a 'Run In' sample to test field procedures. The Run In sample was selected alongside the Year 1 sample using the same methods and was subsequently incorporated into the Year 1 data. The Run In sample was carried out in February and March 2008, therefore an additional set of weights were needed to correct for the unequal distribution of the sample across months in Year 1. The Run In sample weights ensure the full sample (Years 1 to 4 plus the Run In) is seasonally representative.

9.4 Individual non-response weight

The individual non-response weight was generated for the analysis of fully responding individuals; individuals who responded to the individual interview and completed at least three food diary days. The individual non-response weight was generated using calibration weighting methods. An iterative procedure is used to adjust a starting weight until the distribution of the (weighted) sample matches that of the population for a set of key variables. The adjustment keeps the values of the final weights as close as possible to those of the initial weights, which ensures the properties of the initial weights are retained in the final calibrated weights. The initial weights were a set of selection weights. These selection weights corrected for the unequal selection probabilities that were built into the sample design; for the random selection of individuals in households where more than one person was eligible and for the random selection of catering units and dwelling units at multi-unit addresses.

The key variables used to create the individual weight were: age (grouped) by sex and Government Office Region (GOR). The population figures used were taken from the mid-year population estimates.¹⁴ As there are now four years of NDNS RP data, the average population of

the last four years was used. This was generated using the four most recent years of population data (2008-2011).

The aim of the calibration weighting was to reduce non-response bias resulting from differential non-response at the household and individual interview. The calibration weights generated were re-scaled so that the sum of the weights equalled the number of participating individuals; these are the final individual weights (wti_Y1234). Thus the final individual weights adjust for dwelling unit, catering unit and individual selection, the Run In sample, and for the age/sex and regional profiles of participating individuals. This weight should be used for any analyses of interview and food data in the combined Years 1-4 data.

9.5 Nurse interview non-response weight

Participants who completed three or four food diary days (i.e. those deemed fully productive) were asked to consent to a nurse visit. Approximately three quarters of these participants went on to do a nurse interview. Non-response weights were generated to adjust for differences between participants and non-participants to the nurse visit.

There is a small difference in the approach used to create the nurse weights. In previous years the weights were generated in a two-step process; model then calibration. For the Years 1-4 weights the calibration step was dropped as the model produced weights that made the weighted profile of the nurse visit match the weighted profile of all individuals very closely. The modelling step, however, remained the same; a logistic regression was used to model the relationship between response to the nurse interview (coded into a binary outcome variable) and a set of predictor variables (socio-demographic, participant and household/catering unit characteristics collected during the interview). Adults and children were modelled separately as response behaviour can vary between the two groups. The weights from the two models were then combined into a single weight.

The model generated a predicted probability for each participant. This is the probability the participant would take part in the nurse interview, given the characteristics of the individual and the household/catering unit. Participants with characteristics associated with non-response were under-represented in the nurse sample and therefore receive a low predicted probability. These predicted probabilities were then used to generate a set of non-response weights; participants with a low predicted probability got a larger weight, increasing their representation in the sample.

As before, the nurse weights were re-scaled so that the sum of the combined adult and child weights equalled the number of participants who had a nurse visit. These are the final nurse weights for the sample (wtn_Y1234) and adjust for unequal selection, non-response to the household/MFP and individual interviews and non-response to the nurse visit. The nurse weights should be used for all analyses of nurse level data.

9.6 RPAQ non-response weight

All individuals aged 16 years and over were asked to record their physical activity over the previous seven days in a self-completion booklet (the RPAQ).

Response behaviour was modelled using a logistic regression. The same set of predictor variables used to model non-response to the nurse visit was used to model non-response to RPAQ, namely, socio-demographic, participant and household/catering unit characteristics collected during the individual interview. The 16 to 18 year olds were modelled with the adult respondents as children under the age of 16 were not asked to fill in the self-completion booklet.

The RPAQ weights were re-scaled so that the sum of the combined adult and child weights equalled the number of participants who had completed RPAQ. These are the final RPAQ weights for the sample (wtr_Y1234) and adjust for unequal selection, non-response to the household/MFP and individual interviews and non-response to RPAQ.

Note that response to RPAQ and the nurse visit was not hierarchical; it was possible for a respondent to complete the RPAQ section but not the nurse visit, and vice versa.

9.7 ActiGraph non-response weight

In Year 1, all children aged 4 to 10 years were asked to wear an ActiGraph. In Years 2-4, the eligibility criteria were extended to include children aged 11 to 15 years.

The ActiGraph weights are intended to make the ActiGraph sample representative of all eligible children in the population. They were generated using a logistic regression model. The predictor variables tested were a set of socio-demographic participant and household characteristics collected from the individual interview, although only age, sex and region were in the final models. Predicted probabilities were generated for each participant and used to generate a set of non-

response weights. Participants with a low predicted probability received a larger weight, increasing their representation in the sample.

The non-response weights from the model were combined with the final interview weights to give the final ActiGraph weights (the final interview weights incorporate the selection weights and weights for non-response to the individual questionnaire). The weights were combined and then scaled, so the mean combined weight was equal to one and the weighted sample size matched the unweighted sample size.

Note that response to the ActiGraph and the nurse visit was not hierarchical; it was possible for a child to wear an ActiGraph but not go forward to the nurse visit, and vice versa.

9.8 Blood sample non-response weights

An additional set of weights was generated to correct for differential non-response to giving a blood sample. Non-response, whether due to refusal or inability to give a blood sample, will cause the blood data to be biased if there are systematic differences between individuals that provide a blood sample and individuals that do not. Blood samples were taken during the nurse visit. Only participants who fulfilled certain eligibility criteria were asked whether they would be prepared to give a blood sample. Participants were ineligible if they:

- had a clotting or bleeding disorder (e.g. conditions such as haemophilia and low platelets (thrombocytopenia))
- had **ever** had a fit
- were **currently** on anticoagulant drugs, e.g. Warfarin therapy
- had volunteered information that they are HIV or Hepatitis B or C positive

The blood weights were generated using a logistic regression model. Adults and children were modeled separately. The non-response weights from the model were combined with the final nurse weights to give the final blood weights for adults and children (wtb_Y1234). These weights adjust for non-response to the blood sample, non-response to the nurse visit, non-response to the individual questionnaire and unequal selection probabilities. The weights were scaled, so the mean weight equalled one and the weighted sample size matched the unweighted sample size. These weights should be used for any analysis of blood sample data.

9.9 24-hour urine sample non-response weights

The 24-hour urine sample data will be biased if systematic differences between individuals that do and do not provide a complete urine sample are not corrected for. All individuals aged four years and over, with the exception of children still in nappies, were asked by the nurse at stage 2 to provide a 24-hour urine collection.

The analysis needed to exclude, as far as possible, all individuals with incomplete collections without introducing significant bias. Sample completeness was determined by the amount of PABA excretion and whether the respondent reported any missed collections. Individuals who provided an incomplete urine sample were counted as non-responders.

Two different definitions of completeness have been used in this report. The first definition of completeness applies the same criteria as those used for adults in the 2011 Assessment of dietary sodium in adults (aged 19 to 64 years).¹⁵ A collection was deemed to be complete if either the levels of PABA excretion were sufficiently high or (where the individual had declined or failed to take the full PABA requirement) the individual claimed it to be complete. This definition is referred to as the 'standard adult criteria' which were used to identify responding individuals for the first set of weights and applied to all individuals aged four years and over.

The second definition was used for children aged 4 to 10 years. By this definition, children had given a complete urine collection if they reported no missed urine over the required time period. This definition is referred to as 'complete by claim' and were used for the second set of weights to identify responding children aged 4 to 10 years. For all other participants (i.e. those aged 11 years and over) the 'standard adult criteria' was again used for the second set of weights. Hence, the two sets of weights are identical for participants aged 11 years and over.

As children aged 4 to 10 years can be complete by two different criteria, the weighting variables have been generated to reflect this and hence the use of either weighting variable will not provide a full sample of complete cases (i.e. complete by any criterion).

The eligibility criteria meant that participants who provided a usable 24-hour urine sample were weighted to match the eligible nurse participants (i.e. those who were visited by a nurse and were eligible to provide a 24-hour urine sample). It can be assumed that the eligible nurse participants (weighted by the nurse weight) are representative of all eligible persons in the population, since the nurse weights make the full nurse sample representative of the population. The final 24-hour urine

weights therefore make the 24-hour urine sample participants representative of all eligible persons in the population. This assumption is made because there are no available estimates of the actual eligible population (i.e. the population providing a 24-hour urine sample).

24-hour urine collection weights were generated using logistic regression models. Adults and children were modelled separately. Adults were modelled once using an outcome code based on the 'standard adult criteria'. Children were modelled twice: firstly using an outcome based on 'standard adult criteria' and secondly using an outcome that was based on either 'complete by claim' for children aged 4 to 10 years or the 'standard adult criteria' (again) for children aged 11 to 18 years.

The non-response weights from the models were combined with the final nurse weights to give the final 24-hour urine sample weights (the final nurse weights incorporate the selection weights, weights for non-response to the individual questionnaire and weights for non-response to the nurse visit). Two sets of weights are produced: `wtu_Y1234v1` is based on the 'standard adult criteria' for all individuals aged four years and over, `wtu_Y1234v2` is based on complete by claim for individuals aged 4 to 10 years and the 'standard adult criteria' for individuals aged 11 to 15 years. Either weight can be used for analysis of adults, they will give the same weighted frequencies, although weighted totals will be very slightly different due to scaling.

9.10 Selecting the appropriate weight variable

Any analyses that incorporate information from more than one stage should use the weight from the 'last' stage. For example, a cross tabulation using a variable from the nurse visit and a variable from the individual questionnaire should use the nurse weights (i.e. the weights from the latter stage of the survey), similarly, a cross tabulation of blood sample data and data collected at the nurse visit should use the blood weight.

For a more detailed description of the weighting process see Appendix B of the NDNS RP Years 1 to 4 report.

Appendix A UK Nutrient Databank

A.1 Introduction to the UK Nutrient Databank

The food level dataset provides nutrients for the foods in amounts as consumed whilst the UK Nutrient Databank dataset provides nutrient information per 100g.

The UK Nutrient Databank (databank) contains extensive information on the nutrient content of foods commonly consumed in the United Kingdom (UK) and enables nutrient intakes to be calculated from consumption data. The databank was originally developed by the Ministry of Agriculture, Fisheries and Food (MAFF) for the 1990 Dietary and Nutritional Survey of British Adults and subsequently updated for the series of NDNS surveys between 1995 and 2000.¹⁶ Ownership of the databank transferred to the FSA where it was updated for the NDNS of adults aged 19 to 64 years,¹⁷ the Low Income Diet and Nutrition Survey (LIDNS),¹⁸ and prior to commencing the NDNS RP in 2008.¹⁹ In 2010 responsibility for the NDNS RP and ownership of the databank transferred to the Department of Health (DH). In April 2013, responsibility transferred to the Department's Executive Agency, Public Health England (PHE).

Data in the databank (managed by PHE and MRC EWL) is largely the same as that in the published UK food composition tables²⁰ but includes a larger range of processed foods and composite dishes and no missing values. The nutrient data assigned to foods originate from three main sources:

- Ongoing programme of nutrient analysis led by DH
- Food manufacturers and information from food labels
- Recipe calculations for homemade dishes, and some manufactured products

In order to estimate nutrient intakes, values are assigned for all nutrients in all foods. Where reliable information is not available for some nutrients, data were obtained by extrapolating estimates from similar foods. All data were carefully evaluated before being incorporated into the Nutrient Databank.

In the NDNS RP it is essential that the databank is up-to-date and, as far as possible, reflects the nutrient composition of the food supply for each year reported. Hence a programme of updates and revisions is a continuing aspect of the RP, with a yearly update carried out by PHE.²¹ Each year of the RP is coded separately using a contemporaneous version of the databank. Updating of the

databank includes the addition of new foods as well as revision of nutrient composition of existing foods, either at food group level following a programme of reanalysis, or to take account of reformulation by manufacturers and changes in fortification practices. As changes in the databank are partly driven by the availability of new analytical data, new data are produced only occasionally so a gradual change in the nutrient content of the food supply may appear as a step change in the nutrient databank with the same foods having a different composition for some nutrients in one year of the RP compared to the next. It is important to utilise contemporaneous food composition data to produce accurate assessments of dietary intake.

A.2 Description of the Nutrient Databank fields and variables

A.2.1 Food name and food code

All foods are assigned a food code and a clear name is given to describe the code. The name may include specific qualities of the food, informing the user it is a fortified food, or from a specific brand. The food codes in the databank cannot be linked directly to foods in McCance and Widdowson's Composition of Foods.²²

A.2.2 Food code markers

Each food code is assigned as an 'F' or 'R' code. 'F' indicates that the code exists as an atomic food and 'R' denotes a recipe code usually comprising 2 or more food codes. Details of the recipes held in the 'R' codes are not provided in this dataset.

A.2.3 Sub food group name and code

A full breakdown of the NDNS food groups is provided in Appendix R of the NDNS RP Years 1 to 4 report.²³ Each food code is assigned to a subsidiary (sub) food group, expressed as an integer with an alphabetical suffix, which is a food group level of greater detail than the main food groups. The databank displays the sub food group name as well as the code.

A.2.4 Water and vitamin loss

Recipe 'R' food codes may have water and vitamin loss assigned, which provides a correction to the nutrient data for associated cooking losses.

A.2.5 Maximum weight

This is an edit check field to assist users of the nutrient data to pick up any coding errors in portion sizes. Amending the maximum weight will not impact the nutrient values in the databank. Maximum weights are set at sub food group level and do not usually have to be adjusted for each code. Food codes that would not appear in dietary assessment records, such as raw meat, have a maximum weight set at 1.

A.2.6 Base and unit (F codes only)

These fields describe how the nutrient data are expressed. For most foods the nutrients are entered per 100 grams (base = 100, unit = grams). For vitamin and mineral supplements, nutrients are entered per tablet, capsule, teaspoon or drop (base = 1, unit = tablet, capsule, teaspoon or drop). The base and unit of a food code must match the form in which items are recorded in the dietary assessment method.

A.2.7 Dilution

The majority of food codes have a dilution of 1. Concentrated soft drinks and dried products that are made up have a dilution factor greater than 1. For example, if a concentrated drink is usually made up 1 part concentrate to 4 parts water the dilution factor would be 5. This provides the user with additional detail regarding the food in its un-concentrated form; however this is not used in calculating nutrient intakes.

A.2.8 Edible portion

The majority of food codes have an edible portion of 1, however this will be less than 1 for foods that include waste, such as meat weighed with bones.

A.2.9 Comments and descriptions

The comments box provides details on when the food code was created, including any updates and the data source. The description box provides details on the number and brand names of products included in calculating the nutrient information, or the food codes used in collating the recipe. Some food codes (particularly dietary supplements) contain nutritional compounds not measured in the nutrient databank (e.g. vitamin K or lutein), and these are also described in the description field.

A.2.10 Nutrition data

Each food code in the databank has a value assigned for 54 nutrients, including energy, provided in specific units of measurement (see table A.1). Additionally, to ensure accurate reporting of specific food types in NDNS, each food code present in the databank has been disaggregated into 28 specific food components (see table A.2).²⁴ Nutrient values reported as 'trace' are assumed as zero in the nutrient databank. Some data will have been rounded into the appropriate number of decimal places.

A.3 Quality assurance

The databank is designed to perform automated quality assurance checks to nutrient data,²⁵ when food codes are added or updated:

Kcal	$(\text{protein} \times 4) + (\text{fat} \times 9) + (\text{carbohydrate} \times 3.75) + (\text{alcohol} \times 7)$
kJ	$(\text{protein} \times 17) + (\text{fat} \times 37) + (\text{carbohydrate} \times 16) + (\text{alcohol} \times 29)$
Carbohydrate	total sugars + starch
Total sugars	sum of all individual sugars
Total sugars	intrinsic + non milk extrinsic sugars
Total iron	haem iron + non haem iron
Total carotene	$\beta\text{-carotene} + (\alpha\text{-carotene} \times 0.5) + (\beta\text{-cryptoxanthin} \times 0.5)$
Vitamin A	Retinol + (total carotene/6)
Total N	Protein/Nitrogen Conversion Factor (NCF)
Total fat	should be \geq the sum of fatty acids
Fatty acids	$0.6 \times \text{total fat}$ is \leq the sum of the fatty acids.

A.4 Changes to the databank between Years 1 to 4

Year 2 contains a food group (n=149) addition of 60R 1% fat milk. Year 3 contains the food group (n=151) additions of 61R Smoothies 100% fruit and/or juice, and 14B Cheddar cheese.

Year 4 contains a food group (n=153) change of 54A 'Cod liver oil and other fish oils' split into 54N 'Cod liver oil and other fish oils (incl vits A,D,E)' and 54P 'Multivitamin and/or minerals with omega 3'. Along with a change of 54C 'Single vits/minerals not folic, Fe, Ca' which is split into 54L 'Vitamin C only' and 54M 'Single vits or mins not folic acid, Fe, Ca, Vit C'.

Table A.1 Nutrient fields available in the databank

Nutrient (unit of measure)		
Water (g)	Other sugars [^] (g)	Vitamin B6 (mg)
Total Nitrogen (g)	Saturated fatty acids (g)	Vitamin B12 (µg)
Nitrogen conversion factor	Cis- monounsaturated fatty acids (g)	Folate (µg)
Protein (g)	Cis-n3 fatty acids (g)	Pantothenic acid (mg)
Fat (g)	Cis-n6 fatty acids (g)	Biotin (µg)
Carbohydrate (g) [^]	Total trans fatty acids (g)	Sodium (mg)
Energy (kcal)	Cholesterol (mg)	Potassium (mg)
Energy (kJ)	Retinol (µg)+	Calcium (mg)
Alcohol (g)	Total carotene (µg)+	Magnesium (mg)
Englyst Fibre (NSP, g)	Alpha-carotene (µg)+	Phosphorus (mg)
Starch (g)	Beta-carotene (µg)+	Iron (mg)
Total sugars (g)	Beta cryptoxanthin (µg)+	Haem iron [^] (mg)
Non-milk extrinsic sugars* (NMES, g)	Vitamin A (retinol equivalents, µg)+	Non-haem iron (mg)
Intrinsic and milk sugars** (IMS, g)	Vitamin D (µg)	Copper (mg)
Glucose (g)	Thiamin (mg)	Zinc (mg)
Fructose (g)	Riboflavin (mg)	Chloride (mg)
Sucrose (g)	Niacin equivalents (mg)	Iodine (µg)
Maltose (g)	Vitamin C (mg)	Manganese (mg)
Lactose (g)	Vitamin E (mg)	Selenium (µg)

[^] Carbohydrate is expressed as monosaccharide equivalents. Other sugars include oligosaccharides, where data is available on their levels. Haem iron is calculated as 40% of the iron in fish and meat.

* Includes all sugars in fruit juices, table sugar, honey, sucrose, glucose and glucose syrups added to foods + 50% of the sugars in canned, stewed, dried or preserved fruits. Non-milk extrinsic sugars cannot be determined by chemical analysis so values are estimated based on available information on the types and sources of sugar in the food

** Includes all sugars in fresh fruit and vegetables + 50% of the sugars in canned, stewed, dried or preserved fruits + lactose in milk and milk products.

+ Vitamin A retinol equivalent is calculated as Retinol + (total carotene/6). Where, Total carotene is β -carotene + ($\frac{1}{2}$ α -carotene) + ($\frac{1}{2}$ β -cryptoxanthin). Fortified sources of vitamin A are entered as retinol in the nutrient databank.

Table A.2 Disaggregation categories used in the NDNS RP

Disaggregation category (g)	
Fruit (Fresh/Canned Fruit)	Other Red Meat
Dried Fruit	Burgers (Burgers and grillsteaks)
Fruit Juice	Sausages
Smoothie Fruit	Offal
Tomatoes	Poultry
Tomato puree	Processed Poultry
Brassicaceae	Game Birds
Yellow/red/Green (Yellow, Red & Green Leafy Vegetables)	White Fish
Other Vegetables	Oily Fish
Beans (Beans and pulses)	Canned Tuna
Nuts	Shellfish
Beef	Cottage cheese
Lamb	Cheddar Cheese
Pork	Other Cheese
Processed Red Meat	

Appendix B Doubly labelled water (DLW)

Doubly labelled water has been used to derive energy expenditure in a subsample of participants in Years 1 and 3 of the NDNS RP. The information below provides background to how the variables in the dataset have been calculated.

B.1 Total Body Water (L)

From the assumption of a mono-compartmental model of water distribution with elimination following first-order kinetics it follows that the normalised enrichments should decay exponentially with time. Derivation of kinetic parameters from standardised mass spectrometric data was calculated and then transformed by taking the logarithm.

Linear regression was used to find the kinetic parameters N_H , k_H , N_O and k_O , where the subscripts H and O denote hydrogen and oxygen isotopes and N and k are the intercept (body water pool size) and rate constant respectively.

This gives the following equations:

$$\begin{aligned} \ln\{\Delta_H(t)\} &= \ln\left\{\frac{1}{N_H}\right\} - k_H t \\ \ln\{\Delta_O(t)\} &= \ln\left\{\frac{1}{N_O}\right\} - k_O t \end{aligned}$$

This then allowed calculation of total body water (TBW_L), from which body composition can be derived.

$$TBW = \left(\frac{\frac{N_H}{1.04} + \frac{N_O}{1.01}}{2} \right) \left(\frac{18.02}{1000} \right)$$

Where TBW is calculated in kg, the subscripts H and O denote hydrogen and oxygen isotopes and both N_H , and N_O are in moles.

NOTE N_H and N_O are corrected for non-aqueous exchange, with the factors 1.04 and 1.01 respectively.

B.2 Lean Body Mass (kg) and Fat Mass (kg)

Lean body mass (LBM_kg) and fat mass (FM_kg) in kg, was then derived by assuming that fat mass has no associated water and that lean mass is 73% hydrated.

$$LBM = \frac{TBW}{0.73}$$

Fat mass (FM) was calculated by subtracting LBM from body weight.

B.3 Basal Metabolic Rate (according to Schofield's equations, using age, sex, height and weight)

As total energy expenditure (TEE) is highly dependent on age and on the level of physical activity. It can be considered to have two contributing terms: the basal metabolic rate (BMR_kj_day_1), which is the energy requirement to maintain life, and the energy, expended in activity (AEE). The BMR was calculated from predictive equations, according to the following equations:

Age Range	BMR (MJday ⁻¹)	
Under 3 years	Female	$0.068W + 4.281H - 1.730$
	Male	$0.001W + 6.349H - 2.584$
3-10 years	Female	$0.071W + 0.677H + 1.553$
	Male	$0.082W + 0.545H + 1.736$
10-18 years	Female	$0.035W + 1.948H + 0.837$
	Male	$0.068W + 0.574H + 2.157$
18-30 years	Female	$0.057W + 1.184H + 0.411$
	Male	$0.063W - 0.042H + 2.953$
30-60 years	Female	$0.034W + 0.006H + 3.530$
	Male	$0.048W - 0.011H + 3.670$
Over 60 years	Female	$0.033W + 1.917H + 0.074$
	Male	$0.038W + 4.068H - 3.491$

Where W is the subject's weight (kg), and H height (m).

B.4 Total Energy Expenditure (kJ) (calculated according to Schoeller's methods)

The following equation was used to calculate carbon dioxide production. Total energy expenditure was calculated from the CO₂ production rate assuming that 12% of total energy is derived from protein oxidation.

$$F_{CO_2} = \left(\frac{N_O k_O - N_H k_H}{2f_3 + 2.1(f_2 - f_1)} \right) \quad \text{Where } f_1 = 0.941, \quad f_2 = 0.991, \quad \text{and } f_3 = 1.037 \quad \text{are}$$

fractionation factors for 2H leaving the body as water vapour, 18O leaving via the same route, and the exchange of 18O between carbon dioxide and water respectively. This is discussed fully by Coward and Cole.

$$TEE = 22.4(15.480/RQ + 5.550)F_{CO_2} \quad \text{with the respiratory quotient } RQ \text{ taken as 0.85}$$

B.5 Activity energy expenditure (AEE) and physical activity energy expenditure (PAEE)

AEE was calculated as follows:

$$AEE = TEE - BMR$$

Diet induced thermogenesis was estimated as 10% of the total energy level, and on this basis physical activity energy expenditure (PAEE) was calculated as follows:

$$PAEE = 0.9TEE - BMR$$

B.6 Physical activity level (PAL)

PAL is the ratio of TEE and BMR and was calculated as follows:

$$PAL = \frac{TEE}{BMR}$$

PAL values usually range between 1.2 and 2.4, corresponding to bedbound subjects at the low end and those engaged in strenuous work or sports activity at the high end.

¹ <https://www.gov.uk/government/organisations/public-health-england>

² <https://www.food.gov.uk/>

³ Responsibility for nutrition policy in England and Wales transferred from FSA to Health Departments in 2010. Management of NDNS also transferred to the Department of Health in England at that time. From 1 April 2013, responsibility for the survey transferred to the Department of Health's Executive Agency, Public

Health England (PHE).

⁴ Additional recruitment in the devolved countries is funded by Government bodies in Scotland, Wales and Northern Ireland.

⁵ <http://www.natcen.ac.uk/>

⁶ <http://www.mrc-ewl.cam.ac.uk/>

⁷ <http://www.ucl.ac.uk/>

⁸ <http://www.nisra.gov.uk/>

⁹ Ashwell M, Barlow S, Gibson S, Harris C (2006) National Diet and Nutrition Surveys: the British experience. Public Health Nutrition 9(4) 523-530.

¹⁰ Department of Health Healthy Lives, Healthy People: Our Strategy for public health in England White Paper
http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_121941
(accessed 22/04/2014).

¹¹ Run In respondents from Year 1 have a serial number starting with a 9.

¹² Complete by standard criteria: Samples were deemed to be complete and included in the analysis if:

- i) In cases who reported taking 3 PABA tablets, a PABA recovery was between 85-119% by colorimetry or 70-104% by HPLC
- ii) In cases who reported taking <3 PABA tablets or who declined to take PABA, no missed collections were reported and the reported collection period was between 23-25 hours.

¹³ Complete by claim only criterion: Samples were deemed to be complete and included in the analysis if no missing collections were reported and if the reported collection period was 23-25 hours. PABA was not taken into account.

¹⁴ Produced for England and Wales by ONS, for Scotland by GROS and for Northern Ireland by NISRA.

¹⁵ National Diet and Nutrition Survey - Assessment of dietary sodium in adults (aged 19 to 64 years) in England, 2011. Katharine Sadler et al [Online]. Available:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/127916/Sodium-Survey-England-2011_Text_to-DH_FINAL1.pdf (accessed 15/04/2013).

¹⁶ Smithers, G. (1993) MAFF's Nutrient Databank. Nutrition and Food Science; 2; 16-1.

¹⁷ Henderson L, Gregory J, Swan G. National Diet and Nutrition Survey: adults aged 19 to 64 years. Volume 1: Types and quantities of food consumed. London: TSO, 2002.

¹⁸ Nelson M, Erens B, Bates B, Church S, Boshier T. 2007. Low Income Diet and Nutrition Survey. Volume 2. Food Consumption. Nutrient intake. London, The Stationery Office.

¹⁹ National Diet and Nutrition Survey: Results from Years 1, 2 and 3 (combined) of the Rolling Programme 2008/09 – 2010/11[online] <http://transparency.dh.gov.uk/2012/07/25/ndns-3-years-report/>

²⁰ The summary edition and related supplements are referenced and located as follows: Composition of Foods Series, Sixth Edition and Supplements as cited in Bates B, Lennox A, Bates C & Swan G. National Diet and Nutrition Survey: Headline results from Years 1 and 2 (combined) of the Rolling Programme (2008/9 – 2009/10) http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsStatistics/DH_128166

(accessed 20/04/12).

²¹ Previously FSA or DH when ownership of the Nutrient Databank lay with them.

²² McCance and Widdowson's The Composition of Foods integrated dataset [online]
<http://tna.europarchive.org/20110116113217/http://www.food.gov.uk/science/dietarysurveys/dietsurveys/>

²³ National Diet and Nutrition Survey: Appendix R Main and subsidiary food groups [online]
<https://www.gov.uk/government/publications/national-diet-and-nutrition-survey-results-from-years-1-to-4-combined-of-the-rolling-programme-for-2008-and-2009-to-2011-and-2012>.

²⁴ Fitt E, Mak TN, Stephen AM, Prynne C, Roberts C, Swan G & Farron-Wilson M. (2010) Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. Eur J Clin Nutr 64: S32-S36.

²⁵ Results from these calculations may not match the final values exactly (especially when comparing against analytical data) as a small margin of difference is acceptable in food composition.